

Fireball Loading and the Blast Wave Model of Gamma Ray Bursts

Charles D. Dermer¹, James Chiang^{1,2}, and Markus Böttcher^{1,3}

ABSTRACT

A simple function for the spectral power $P(\epsilon, t) \equiv \nu L(\nu)$ is proposed to model, with 9 parameters, the spectral and temporal evolution of the observed nonthermal synchrotron power flux from GRBs in the blast wave model. Here $\epsilon = h\nu/m_e c^2$ is the observed dimensionless photon energy and t is the observing time. Assumptions and an issue of lack of self-consistency are spelled out. The spectra are found to be most sensitive to the baryon loading, expressed in terms of the initial bulk Lorentz factor Γ_0 , and an equipartition term q which is assumed to be constant in time and independent of Γ_0 . Expressions are given for the peak spectral power $P_p(t) = P(\epsilon_p, t)$ at the photon energy $\epsilon = \epsilon_p(t)$ of the spectral power peak. A general rule is that the total fireball particle kinetic energy $E_0 \sim \Pi_0 t_d$, where $t_d \propto \Gamma_0^{-8/3}$ is the deceleration time scale and $\Pi_0 \equiv P(\epsilon_p, t_d) \propto \Gamma_0^{8/3}$ is the maximum measured bolometric power output in radiation, during which it is carried primarily by photons with energy $\mathcal{E}_0 = \epsilon_p(t_d) \propto q \Gamma_0^4$.

This rule governs the general behavior of fireballs with different baryon loading. Clean fireballs with small baryon loading ($\Gamma_0 \gg 300$) are intense, subsecond, medium-to-high energy γ -ray events, and are difficult to detect because of deadtime and sensitivity limitations of previous γ -ray detectors such as EGRET on *CGRO*. Dirty fireballs with large baryon loading ($\Gamma_0 \ll 300$) produce transient emissions which are longer lasting and most luminous at X-ray energies and below, but these events are lost behind the glow of the X-ray and lower-energy background radiations except for rare serendipitous detections by pointed instruments. The correlation between hardness and duration of loaded GRB fireballs ($100 \lesssim \Gamma_0 \lesssim 1000$) follows from this rule.

1. Introduction

The cosmological origin of GRBs has been established as a result of follow-up observations of fading X-ray counterparts to GRB sources discovered with the *Beppo-SAX* mission (e.g., Costa et al. 1997; Metzger et al. 1997; van Paradijs et al. 1997; Djorgovski et al. 1997; Kulkarni et al. 1998a). The time profiles of the X-ray afterglow light curves are generally well fit by power laws with temporal indices χ in the range $1.1 \lesssim \chi \lesssim 1.5$ where, at a fixed observing frequency, the flux $\phi \propto t^{-\chi}$ (e.g, Costa et al. 1997; Feroci et al. 1998). Only the X-ray afterglow behavior of GRB 970508 is not well fit with a single power law, as its X-ray flux displays a peak between 1-2 days after the GRB. Its overall X-ray behavior from 30 seconds to 1 week following the burst event does, however, follow a power-law decay with $\chi = 1.1$ (Piro et al. 1998a). The light curves of GRB counterparts detected at optical and radio frequencies also generally display power-law behavior with a similar range of temporal indices (e.g., Sahu et al. 1997; Galama et al. 1998; but note Groot et al. 1998; Frail 1998).

¹E. O. Hulburt Center for Space Research, Code 7653, Naval Research Laboratory, Washington, DC 20375-5352

²NRL/NRC Resident Research Associate

³Department of Space Physics and Astronomy, Rice University, Houston, TX 77005-1892

The long wavelength afterglows of GRBs are most simply explained with the fireball/blast-wave model, and were predicted prior to their discovery (Paczynski & Rhoads 1993; Mészáros & Rees 1997). In this model, the impulsive release of a large amount of energy in a small volume is transformed into relativistic plasma whose baryon-loading parameter, expressed in terms of the initial bulk Lorentz factor Γ_0 of the fireball after it has become optically thin, obtains values much greater than unity (Blandford & McKee 1976; Cavallo & Rees 1978; Shemi & Piran 1990; Rees & Mészáros 1992; Piran & Shemi 1993). The dominant radiation mechanism producing the prompt GRB emission and radio through X-ray afterglows is nonthermal synchrotron emission (Mészáros & Rees 1993a; Katz 1994; Tavani 1996; Cohen et al. 1997). The power-law behavior of the long wavelength afterglows is a consequence of the deceleration of the blast wave as it sweeps up material from the ambient medium (Wijers, Rees, & Mészáros 1997; Waxman 1997a; Vietri 1997a; Dermer & Chiang 1998). The blast-wave model also provides an explanation for the > 100 MeV emission observed with the EGRET instrument from bright BATSE bursts (Dingus 1995; Catelli, Dingus, & Schneid 1996; Hurley et al. 1994) as due either to synchrotron self-Compton (SSC) emission (Mészáros, Rees, & Papathanassiou 1994; Chiang & Dermer 1998) or to the nonthermal proton synchrotron radiation from hadrons accelerated in the relativistic blast wave (Vietri 1997b; Böttcher & Dermer 1998).

In this paper, we propose a simple function to describe the observed prompt and delayed emissions from a decelerating blast wave in the fireball/blast-wave model. We use this function to perform a parameter study, and find that the temporal and spectral behavior of the blast-wave emissions are robust to orders-of-magnitude changes in all parameter values except for the baryon-loading parameter Γ_0 and an equipartition parameter q . If q is not strongly dependent on Γ_0 , as quantified below, then the observability of a fireball is most strongly determined by the value of the baryon-loading factor Γ_0 . Clean fireballs with small baryon loading ($\Gamma_0 \gg 300$) produce intense, subsecond, medium-to-high energy ($\gg 10$ MeV) γ -ray events, and heavily-loaded dirty fireballs ($\Gamma_0 \ll 300$) produce longer duration transients which are most luminous at X-ray energies and below. Limitations of earlier telescopes may have prevented the discovery of clean fireballs, though some previous serendipitous detections of X-ray transients might represent members of the dirty fireball class.

In Section 2, we introduce a function to model the temporally-evolving nonthermal synchrotron emission produced by relativistic blast waves which decelerate and radiate by sweeping up particles from the circumburst medium (CBM). The terms in this function are derived from basic blast-wave physics. Differences in the emission properties of blast waves due to changes in parameter values are examined in Section 3. In particular, the strong-dependence of the blast-wave spectra on Γ_0 is noted. In Section 4, we consider the limitations of and requirements for telescopes to be capable of detecting clean and dirty fireballs in view of the emission properties of blast waves with different baryon loading. The predictions of clean and dirty fireballs as new classes of astrophysical transients are discussed in Section 5, where the central results of this study are summarized through a simple rule which governs the overall emission properties of blast waves produced by relativistic fireballs.

2. Spectral Power and Spectral Power Flux of GRBs in the Blast Wave Model

GRB emissions are most conveniently described in terms of their measured νF_ν spectral power fluxes. For simplicity, we consider only uncollimated, spherically expanding blast waves here. The spectral power $P(\epsilon, t) = \nu L_\nu$ is related to the spectral power flux through the relation $P(\epsilon, t) = 4\pi d_L^2 \nu F_\nu$, where d_L is the luminosity distance. Due to the strong Doppler beaming when $\Gamma \gg 1$, most of the observed flux is produced by emission regions located within an angle $\theta \lesssim 1/\Gamma$ of the line-of-sight direction. As the blast

wave decelerates into and through the nonrelativistic regime, the observer receives additional contributions from emission regions at larger angles to the line-of-sight (Panaitescu & Mészáros 1998a; Waxman 1997b; Sari 1998; Chiang & Dermer 1998). This introduces corrections to the above relation between the produced power and observed flux, but this effect is not important when $\Gamma \gg 1$, and is neglected here.

In this limit, we model the temporally-evolving GRB spectrum with the function

$$P(\epsilon, t) \left[\epsilon \times \frac{\text{ergs}}{s - \epsilon} \right] = \frac{(1 + v/\delta) P_p(t)}{[\epsilon/\epsilon_p(t)]^{-v} + (v/\delta)[\epsilon/\epsilon_p(t)]^\delta} \quad (1)$$

where, on a νL_ν plot, the rising slope has index v (*upsilon*), and the descending slope has index δ (*down*). Both v and δ are > 0 . The dimensionless photon energy ϵ is in units of m_e , where particle masses are in energy units. Obviously $P(\epsilon_p, t) = P_p(t)$. Expression (1) applies only to the nonthermal synchrotron portion of the spectrum, in accord with detailed treatments (Chiang & Dermer 1998; Panaitescu & Mészáros 1998b) which show that the SSC contribution to the total power output is generally much weaker than the nonthermal electron synchrotron contribution. The SSC component can still represent the dominant emission process in certain frequency ranges, for example, at $\gg 100$ MeV gamma-ray energies or at soft gamma-ray energies in the afterglow phase when the high-energy flux is at a very low level. At MeV energies and below, however, the SSC component can usually be neglected. Compton processes involving external photons can also be shown to be negligible unless the surrounding radiation fields have extremely large energy densities $\gtrsim 10^4$ eV cm $^{-3}$ (see, e.g., Dermer, Sturmer, & Schlickeiser 1997).

We assume that the density of the CBM surrounding the location of the burst event can be parameterized by a power law in distance from the center of the explosion. Thus

$$n(x) = n_0 x^{-\eta} . \quad (2)$$

The dimensionless spatial coordinate x represents the distance from the explosion site in units of the deceleration radius

$$x_d = \left[\frac{(3 - \eta)E_0}{4\pi n_0 \Gamma_0^2 m_p} \right]^{1/3} , \quad (3)$$

which characterizes the distance at which the blast wave has swept up $\approx 1/\Gamma_0$ times the initial baryonic mass of the fireball (Rees & Mészáros 1992; Mészáros & Rees 1993b).

It is useful to parameterize the evolution of the bulk Lorentz factor of the blast wave by the expression

$$\Gamma(x) = \begin{cases} \Gamma_0 , & \text{if } 0 \leq x < 1; \\ \Gamma_0 x^{-g} , & \text{if } 1 \leq x \leq \Gamma_0^{1/g} . \end{cases} \quad (4)$$

The index g refers to the deceleration regime. If very little of the swept-up energy is radiated over the time scale on which the blast wave decelerates, then the blast wave evolves in the non-radiative (or adiabatic regime) with $g \rightarrow 3/2$ when $\eta = 0$ (Blandford & McKee 1976; Paczyński & Rhoads 1993). This has a self-similar solution independent of Γ_0 when $x \gg 1$. If, on the other hand, the bulk of the swept-up energy is promptly radiated away, then the blast wave evolves in the radiative regime with $g \rightarrow 3$ when $\eta = 0$ (e.g., Mészáros, Rees, & Wijers 1997 and references therein). Regimes intermediate to these two limits have been treated analytically (Dermer & Chiang 1998) and numerically (Chiang & Dermer 1998; Panaitescu & Mészáros 1998c).

Corresponding to x_d is the deceleration time scale in the observer's frame, given by

$$t_d = \frac{(1 + z)x_d}{\Gamma_0^2 c} = \frac{(1 + z)}{c \Gamma_0^{8/3}} \left[\frac{(3 - \eta)E_0}{4\pi n_0 m_p} \right]^{1/3} . \quad (5)$$

Eqs. (3) and (5) hold when $\eta < 3$, and are readily generalized for more complicated radial structure involving, for example, inner and outer boundaries of the CBM, or more general radial distributions with $\eta = \eta(r)$. In the most general case, $\eta = \eta(\mathbf{r})$. A more complicated radial dependence of η than given through eq. (2) would produce additional time-dependent variations in the spectral output of the decelerating blast wave than derived here. For example, short time scale variability in GRB light curves may result from clouds surrounding the GRB source (Dermer & Mitman 1998). The dimensionless spatial coordinate x is related to the observed time t through the expression

$$x = \begin{cases} t/t_d, & \text{if } 0 \leq t < t_d \text{ and } 0 \leq x < 1; \\ [(2g+1)\frac{t}{t_d} - 2g]^{1/(2g+1)}, & \text{if } t_d \leq t \leq \frac{t_d}{(2g+1)} \left(\Gamma_0^{2+g^{-1}} + 2g \right) \text{ and } 1 \leq x \leq \Gamma_0^{1/g}. \end{cases} \quad (6)$$

The existence of a low-energy cutoff in the electron distribution function is reasonable, given that the blast wave sweeps up electrons and protons from the CBM with Lorentz factor Γ in the comoving frame. If an efficient mechanism transfers energy from the protons to the electrons, then the low-energy cutoff of the electron Lorentz factors can reach values as large as

$$\gamma_{e,\min} = \xi_e (m_p/m_e) \Gamma \quad (7)$$

(e.g., Mészáros et al. 1994). The term $\xi_e < 1$ is an electron equipartition factor. Synchrotron radiation from an electron distribution with a low-energy cutoff produces a low-energy spectrum with $\nu = 4/3$ (Katz 1994; Tavani 1996; Cohen et al. 1997).

Fermi processes in the blast wave shock can additionally accelerate a nonthermal component of electrons and protons. If the injection index is steeper than 3, or if the injection index is steeper than 2 and cooling processes are rapid, then the νF_ν spectrum falls with $\delta > 0$. It is assumed here that the values of ν and δ and, by implication, the spectrum of the underlying particle distributions do not evolve with time. This is clearly not the case due to particle cooling through radiative processes (see Chiang & Dermer 1998 for a numerical treatment). These processes must be carefully considered for interpretations of and fitting to afterglow spectra. But these details are not so important for the overall energetics of the blast-wave emissions considered here.

Another crucial uncertainty of blast-wave models is the magnetic field strength H in the comoving blast wave frame. This is commonly given in terms of an “equipartition field” obtained by equating the magnetic field energy density with the nonthermal particle energy density of the swept-up particles downstream of the forward shock. Thus H is parameterized by the expression

$$H(\text{G}) = [32\pi m_p n(x) \xi_H(r/4)]^{1/2} \Gamma(x) = 116 [n(x) \xi_H(r/4)]^{1/2} \Gamma_{300}, \quad (8)$$

where $n(x)$ is the density of the swept-up gas in the stationary frame of the GRB source given by eq. (2), r is the compression ratio, $\Gamma_{300} = \Gamma(x)/300$, and ξ_H is the magnetic-field equipartition parameter. Field generation through dynamo processes could strengthen H with time (see, e.g., Ryu & Vishniac 1991; Mészáros, Laguna, & Rees 1993 and references therein); flux-freezing and reconnection could weaken it (e.g., Mészáros et al. 1994). The choice $\xi_H \sim 1$ does not produce GRB spectral forms similar to those observed, as excessive cooling produces a spectral component with $\nu \cong 0.5$ (Sari, Piran, & Narayan 1998; Chiang & Dermer 1998). Values of $\xi_H \lesssim 10^{-4}$ are required to give good fits to GRB spectra during the prompt phase. Although there is no justification for treating ξ_H as time-independent when $\xi_H \neq 1$, we do so here for simplicity.

In a tangled field with mean field strength H given by eq. (8), the low-energy electron Lorentz-factor cutoff (7) gives a peak in the spectral power at energy $H\gamma_{e,\min}^2/H_{\text{cr}}$ in the comoving frame, where $H_{\text{cr}} = 4.414 \times 10^{13}$ G. To the observer, this energy is boosted and redshifted by a factor $\approx \Gamma/(1+z)$. Thus the observed peak energy ϵ_p at time t is given by

$$\epsilon_p(t) = \mathcal{E}_0[\Gamma(x)/\Gamma_0]^4 x^{-\eta/2} = \mathcal{E}_0 \begin{cases} x^{-\eta/2}, & \text{if } 0 \leq x < 1; \\ x^{-4g-\eta/2}, & \text{if } 1 \leq x < \Gamma_0^{1/g}, \end{cases} \quad (9)$$

where

$$\mathcal{E}_0 = \frac{3.0 \times 10^{-8} n_0^{1/2} q \Gamma_0^4}{(1+z)}, \quad (10)$$

and

$$q \equiv [\xi_H(r/4)]^{1/2} \xi_e^2. \quad (11)$$

The coefficient in eq. (10) is $(m_p/m_e)^2(32\pi m_p)^{1/2}/H_{\text{cr}} = 2.97 \times 10^{-8}$, and $\mathcal{E}_0$ represents the observed photon energy of the peak of the νF_ν spectrum when the blast wave passes through the deceleration radius at $x = 1$.

The power in swept-up particle kinetic energy in the comoving frame is

$$\dot{E}_{\text{ke}} = m_p c B(x) \Gamma(x) [\Gamma(x) - 1] n(x) A(x) \quad (12)$$

(Blandford & McKee 1976), where $A(x) = A_0 x^2 = (4\pi x_d^2) x^2$ is the area of the spherically expanding blast wave and $B(x)c = [1 - \Gamma(x)^{-2}]^{1/2} c$ is its speed. If ζ is defined as the fraction of swept-up energy retained in the comoving frame, then $(1 - \zeta)$ is the fraction that is dissipated by the blast wave shock. Although this dissipation could be in the form of energetic particles which escape directly from the shock, or in the form of bulk kinetic energy which is randomized and stored in the reverse shock to be radiated away later or used to accelerate the blast wave front (see, e.g., Mészáros & Rees 1997; Panaitescu & Mészáros 1998a,c), we assume that essentially all of the dissipation occurs in the form of photon radiation.

As previously discussed, only the emitting region along the line of sight makes a significant contribution to the observed flux in the limit $\Gamma \gg 1$. The observed power is boosted and redshifted by the factor $\Gamma^2/(1+z)^2$ over the radiant power in the comoving frame. Consequently we obtain an expression relating the observed power to the swept-up power, given by

$$\dot{E}_{\text{rad}} \cong \frac{\Gamma^2}{(1+z)^2} (1 - \zeta) \dot{E}_{\text{ke}} \cong (1 - \zeta) m_p c \Gamma^4 n(x) A(x) / (1+z)^2. \quad (13)$$

The spectral power given by eq. (1) satisfies the normalization

$$E_0 = \int_0^\infty dt \dot{E}_{\text{rad}}(t) = \int_0^\infty dt \int_0^\infty d\epsilon \epsilon^{-1} P(\epsilon, t), \quad (14)$$

where E_0 is the total energy in baryons at the end of fireball coasting phase. Thus the radiated power

$$\dot{E}_{\text{rad}} \cong 2(v^{-1} + \delta^{-1}) P_p(t). \quad (15)$$

We use the approach of Dermer & Chiang (1998) to find that in the regime $1 \ll x \ll \zeta \Gamma_0^{1/g}$, the fraction of swept-up energy that is dissipated promptly is given by $(1 - \zeta) = (2g - 3 + \eta)/g$ (see Appendix), and we

assume that this relation holds elsewhere as well. Consequently we derive an expression for the measured bolometric spectral power, given by

$$P_p(t) = \Pi_0 \begin{cases} x^{2-\eta}, & \text{if } 0 \leq x < 1 ; \\ x^{2-\eta-4g}, & \text{if } 1 \leq x < \Gamma_0^{1/g} . \end{cases} \quad (16)$$

The coefficient

$$\Pi_0 = \frac{(2g-3+\eta)m_p c \Gamma_0^4 n_0 A_0}{2g(v^{-1} + \delta^{-1})(1+z)^2} = \frac{c(2g-3+\eta)\Gamma_0^{8/3}}{2g(v^{-1} + \delta^{-1})(1+z)^2} (4\pi m_p n_0)^{1/3} (3-\eta)^{2/3} E_0^{2/3} \quad (17)$$

represents the bolometric luminosity (ergs s⁻¹) at the time t_d of peak power output. We again note that a more complicated expression would hold if η were not constant, thereby rendering ζ dependent upon time.

It is straightforward to determine the time-dependence of $\epsilon_p(t)$ and $P_p(t)$ in different regimes from eqs. (6), (9), and (16). For the parameterized density distribution (2), the peak photon energy $\epsilon_p(t) \propto t^{-\eta/2}$ when $t \leq t_d$, whereas for $t \geq t_d$, $\epsilon_p(t) \propto t^{-(12+\eta)/8}$ and $\propto t^{-(24+\eta)/14}$ in the non-radiative and radiative regimes, respectively. For a uniform medium ($\eta = 0$), $\epsilon_p(t)$ is constant for $t \leq t_d$, whereas for $t \geq t_d$, $\epsilon_p(t) \propto t^{-3/2}$ and $\propto t^{-12/7}$ in the non-radiative and radiative regimes, respectively. The peak spectral power $P_p(t)$ increases $\propto t^{2-\eta}$ for $t \leq t_d$, and $P_p(t) \propto t^{-1-\eta/4}$ and $\propto t^{-(10+\eta)/7}$ in the limiting non-radiative ($g = 3/2$) and radiative ($g = 3$) regimes, respectively, when $t \gg t_d$. For a uniform medium with $\eta = 0$, $P_p(t) \propto t^2$ for $t \leq t_d$, and $P_p(t) \propto t^{-1}$ in the non-radiative regime when $t \geq t_d$ (Sari 1997). In the radiative regime, $P_p(t) \propto t^{-10/7}$ when $t \geq t_d$. From these relations, one can see that the power emitted in lower-energy radiation such as the optical band will always be less than in the higher energy X-ray and gamma-ray bands for declining density profiles (i.e., $\eta > 0$).

One can also easily determine the temporal indices measured at a fixed observing energy ϵ by consulting equations (1), (6), (9), and (16). Let $t_p(\epsilon)$ represent the time when the spectral power flux at ϵ reaches its maximum value. It is found by solving $\epsilon = \epsilon_p(t_p)$ using eqs. (6) and (9), yielding

$$t_p(\epsilon) = \frac{t_d}{2g+1} \left[\left(\frac{\epsilon}{\mathcal{E}_0} \right)^{-\frac{2g+1}{4g+\eta/2}} + 2g \right]. \quad (18)$$

Note that $t_p(\epsilon) \rightarrow t_d$ when $\epsilon \gg \mathcal{E}_0$, whereas $t_p(\epsilon) \gg t_p$ when $\epsilon \ll \mathcal{E}_0$. The displacement of the pulse peaks at lower energies may act alone or in concert with synchrotron cooling (Chiang 1998; Dermer 1998) to produce the observed energy dependence of the GRB pulse width (Fenimore et al. 1995; Piro et al. 1998b). Effect (18) represents a prediction of this model, and will be explored in more detail elsewhere (Dermer, Böttcher, & Chiang 1998).

Two regimes characterize the time-dependence of the flux at high energies $\epsilon \gg \mathcal{E}_0$, namely

$$P(\epsilon, t) = \left(1 + \frac{\delta}{v}\right) \Pi_0 \left(\frac{\epsilon}{\mathcal{E}_0}\right)^{-\delta} \times \begin{cases} (t/t_d)^{2-\eta-\eta\delta/2}, & \text{if } t \ll t_d, \\ [(1+2g)t/t_d]^{[2-\eta(1+\delta/2)-4g(1+\delta)]/(2g+1)}, & \text{if } t \gg t_d. \end{cases} \quad (19)$$

In contrast, three regimes characterize the time-dependence of the flux at energies $\epsilon \ll \mathcal{E}_0$. They are

$$P(\epsilon, t) = \Pi_0 \times \begin{cases} (1+v/\delta)(\frac{\epsilon}{\mathcal{E}_0})^v (t/t_d)^{2-\eta+\eta v/2}, & \text{if } t \ll t_d, \\ (1+v/\delta)(\frac{\epsilon}{\mathcal{E}_0})^v [(1+2g)t/t_d]^{[2-\eta(1-v/2)-4g(1-v)]/(2g+1)}, & \text{if } t_d \ll t \ll t_p(\epsilon), \\ (1+\delta/v)(\frac{\epsilon}{\mathcal{E}_0})^{-\delta} [(1+2g)t/t_d]^{[2-\eta(1+\delta/2)-4g(1+\delta)]/(2g+1)}, & \text{if } t \gg t_p(\epsilon). \end{cases} \quad (20)$$

Note that the late time behaviors in eqs. (19) and (20) are identical.

At times $t > t_d$, two types of behavior occur depending on whether the observations take place at $\epsilon \ll \mathcal{E}_0$ or $\epsilon \gg \mathcal{E}_0$. In the former case, there are two branches in the time profile related to the time $t_p(\epsilon)$ when ϵ_p sweeps through the observing energy ϵ . When $t_d \ll t \ll t_p(\epsilon)$,

$$P(\epsilon, t) \propto t^{\chi_1} \quad \text{where} \quad \chi_1 = [2 - \eta(1 - v/2) - 4g(1 - v)]/(2g + 1). \quad (21)$$

When $t \gg t_p(\epsilon)$, the time profile decays according to the relation

$$P(\epsilon, t) \propto t^{-\chi_2} \quad \text{where} \quad \chi_2 = [4g(1 + \delta) + \eta(1 + \delta/2) - 2]/(2g + 1) \quad (22)$$

In the latter high-energy case ($\epsilon \gg \mathcal{E}_0$), a single power-law decay in the time profile is found for $t \gg t_d$ which follows the behavior given by eq. (22). Table 1 lists the derived temporal slopes χ_1 and χ_2 for various values of v, δ , and g .

3. Model Spectra

Table 2 gives a list of the parameters which, when placed in eqs. (5), (6), (9), (10), (16), and (17), can be used to calculate temporally-evolving model spectra using spectral form (1). Fig. 1a shows such a calculation employing the parameter values listed in Table 2 and called “standard” henceforth. For these parameters, the deceleration time scale $t_d = 58$ s and $\mathcal{E}_0 = 0.4$, giving a duration and a 200 keV νF_ν peak energy early in the event which are characteristic of typical GRBs. The smoothly joined broken power-law form resembles the canonical GRB “Band”-type spectrum (Band et al. 1993). This model burst radiates a peak power $\Pi_0 \cong 3.8 \times 10^{50}$ ergs s⁻¹ at t_d and would be below the BATSE flux threshold at redshift $z = 5$, but would be strongly detected if the GRB source were at $z = 1$.

The spectra in Fig. 1a are plotted at observing times from 1 μ s to 10⁷ s in factor-of-10 increments. The solid lines which overplot the spectra connect the peak energies $\epsilon_p(t)$ and peak luminosities $P_p(t)$ at different observing times. When $t \leq t_d$, $\epsilon_p(t)$ is constant because $\eta = 0$ in this example, whereas it declines at later times according to the relation $\epsilon_p(t) \propto t^{-4g/(2g+1)} \propto t^{-1.6}$, noting eqs. (6) and (9) with $g = 2$. For these same parameters, the peak power output $P_p(t) \propto t^2$ when $t \leq t_d$, but $P_p(t)$ decreases $\propto t^{(2-4g)/(2g+1)} \propto t^{-6/5}$ when $t \gg t_d$, noting eqs. (6) and (16). We see that observations of the temporal decay of the νF_ν peak relate the indices g and η which characterize the radiative regime and density profile of the external medium, respectively. Again, the possible spatial dependence of η complicates any simple interpretation of the measured behavior.

Fig. 1b is a representation of the model data shown in Fig. 1a in the form of time profiles at different observing energies. Light curves are shown at $\epsilon = 10^{-12}, 10^{-9}, 10^{-6}, 10^{-3}, 1, 10^3$, and 10^6 , roughly corresponding to radio, mm, optical, X-ray, MeV, GeV, and TeV frequencies, respectively. The light curves at $t > t_d$ display one or two power-law behaviors depending on whether $\epsilon > \mathcal{E}_0 = 0.4$ or $\epsilon < 0.4$, respectively, for the reasons given in the previous section. For the very late time behavior with $t \gg t_p(\epsilon) > t_d$, the temporal decay index $\chi_2 = [4g(1 + \delta) - 2]/(2g + 1) = 1.52$, in agreement with the behavior shown in Fig. 1b. It is interesting to note that temporal indices in the observed range $1.1 \lesssim \chi \lesssim 1.5$ require flat spectra with $\delta \lesssim 1/3$ when $\eta = 0$ (see Table 1).

The GRB spectral indices v and δ are assumed for simplicity not to evolve with time. Thus spectra at photon energies $\epsilon \ll \epsilon_p$ and $\epsilon \gg \epsilon_p$ rise and fall with the indices $v = 4/3$ and $\delta = 0.2$, respectively, for the

parameters given in Table 2. To examine the effects of different parameter values on the observed spectra, it is therefore sufficient to plot the $P_p(t)-\epsilon_p(t)$ trajectory at various times, as indicated in Fig. 1a by the straight lines. Figs. 2a-e presents a parameter study in this form. The Table 2 standard is shown in each of these figures by the solid lines with open circles at times incremented by factors-of-10 starting from $1 \mu\text{s}$.

Fig. 2a shows the effects of changing the index which characterizes the radiative regime. Here we show two cases with $g = 1.6$ and $g = 2.9$, in addition to the $g = 2$ standard. The case $g = 1.6$ is near the non-radiative limit (if $g = 3/2$, there is no radiation by definition), and $g = 2.9$ is near the fully radiative limit. The weakly radiative blast wave with $g = 1.6$ persists for a longer period of time, though it radiates less power in its early phases. The highly radiative blast wave with $g = 2.9$ is nearly an order-of-magnitude more luminous in its early phases than the $g = 1.6$ case, but its inertia drops more quickly and it therefore has a shorter luminous life than in the less radiative cases.

At this point, a remark concerning a serious issue regarding lack of self-consistency of the parameter list is in order. The type of radiative regime specified by g essentially gives the fraction of swept-up power that is dissipated. If nonthermal synchrotron radiation represents the dominant power drain from the blast wave, as we assume, then a fraction $\approx 2 - g^{-1}(3 - \eta)$ of the incoming swept-up power is promptly transformed into radiation. This places demands on the underlying particle distribution and comoving energy densities which are not considered here, but must be kept in mind in any analytic treatment employing a parameterization of blast-wave dynamics according to an expression like eq. (4). A detailed numerical simulation with a self-consistent treatment of plasmoid/blast-wave dynamics is essential to model spectra accurately (Chiang & Dermer 1998).

Figs. 2b and 2c show the effects of varying the density n_0 and the explosion energy E_0 , respectively, over four orders of magnitude. In higher density environments, a blast wave’s energy is radiated more rapidly and, provided q is the same, in the form of higher energy photons; thus its peak power output obtains larger values than in more dilute environments. Fireballs with larger total energies but with equal baryon-loading persist for longer periods of time because both t_d and the time scale to decelerate to nonrelativistic energies are $\propto E_0^{1/3}$ (see eqs. [5] and [6]). For a fixed baryon loading factor (or constant Γ_0), the overall duration of the GRB is multiplied by a factor $\propto (E_0/n_0)^{1/3}$ as a consequence of the deceleration time scale (see eqs.[3] and [5]). This dependence comes from dimensional analysis in scaling solutions for spherical hydrodynamics, and is present in both the Sedov solution (see, e.g., Lozinskaya 1992) and its relativistic equivalent derived by Blandford & McKee 1976). The peak power output changes to compensate for the different duration; thus $P_p(t_d) \propto E_0^{2/3}$ when E_0 varies and n_0 is held constant, and $P_p(t_d) \propto n_0^{1/3}$ when n_0 varies and E_0 is constant (eq. [17]). This latter dependence is a function of the radiative efficiency, which is assumed to be identical to the factor $(1 - \zeta)$ derived in the Appendix. Consequently, the dimensional dependence of $P_p(t)$ is more complicated than that of t_d . In any case, the durations and powers are not strongly dependent on changes of n_0 and E_0 .

Fig. 2d shows the effect of varying the equipartition term q over four orders-of-magnitude, from $q = 10^{-5}$ to $q = 10^{-1}$. A change in q by a constant factor causes the curves to be displaced horizontally by the same constant factor, since $\epsilon_p(t) \propto q$ but $P_p(t)$ is independent of q .

The effects of different amounts of baryon loading are shown in Fig. 2e. The spectra are found to be extremely sensitive to changes in Γ_0 . The mean photon energy $\mathcal{E}_0 = \epsilon_p(t_d)$ at the time of peak-power output Π_0 is $\propto \Gamma_0^4$. Thus most of the radiation of a clean fireball with $\Gamma_0 = 3000$ is in the form of photons with energies eight orders of magnitude larger than the emission from a dirty fireball with $\Gamma_0 = 30$, provided that q remains constant. The duration of peak power output is just the deceleration timescale t_d which is

$\propto \Gamma_0^{-8/3}$. To compensate for the constant total energy in the fireball, $\Pi_0 \propto \Gamma_0^{8/3}$. Thus the duration of peak power output of the clean fireball case is a factor $\sim 2 \times 10^5$ shorter than that of the dirty fireball case in this example, but the magnitude of the power emitted during this period is a factor of $\sim 2 \times 10^5$ greater.

Clean fireballs produce extremely short pulses of radiation which are carried by very high-energy photons. For example, when $\Gamma_0 = 3000$, the peak power output of $\approx 10^{53}$ ergs is carried by $\gtrsim 1$ GeV photons in a burst lasting $\lesssim 0.1$ s. Again, the total energy remains roughly constant, though it is carried by many fewer, though much higher-energy photons. Its late-time behavior approaches that of fireballs with smaller values of Γ_0 , but with lower power to compensate for the energy radiated during the early phases of the event. For the dirty fireball with $\Gamma_0 = 30$, the peak power output at UV energies lasts for ~ 10 minutes and develops into a broadband radiation pulse which evolves to lower energies at later times. Incidentally, this behavior is found in blazar flares (e.g., Marscher & Gear 1985), though a closer examination must be made to see if this model provides an adequate explanation for such phenomena.

The GRBs which trigger BATSE display a very broad bimodal duration distribution ranging from ~ 30 ms to ≈ 500 seconds (Kouveliotou et al. 1993; Belli 1995). A short, hard component of GRBs is seen with durations ranging from ≈ 30 ms to 1 second, with a mean duration of ≈ 0.2 seconds. A longer duration component ranges from ≈ 2 -500 seconds, with a mean duration of 15 - 30 seconds. This hardness-duration correlation is in accord with the behavior outlined above for blast waves with different baryon-loading, where the class of short, hard bursts are produced by fireballs which are less baryon-loaded than average. A model for GRBs in terms of a distribution of Γ_0 is needed, however, to establish whether these GRBs constitute a separate class, or an extension of the typical loaded fireball in the cleaner limit (Dermer et al. 1998). Different levels of baryon-loading might be expected if GRBs originate from multiple sources, as indicated if the claimed association of GRB 980425 with SN 1998bw is correct (Kulkarni et al. 1998b). For the purposes of this paper, we consider a continuous range of fireball-loading parameters rather than different levels of baryon loading.

Thus we find that the observed spectra from fireball/blast waves, which decelerate and are energized by the process of sweeping up material from the CBM, are most strongly dependent on the baryon-loading parameter Γ_0 and the equipartition parameter q . The term q determines the mean photon energy of the radiant emission through the relation $\mathcal{E}_0 \propto q\Gamma_0^4$ (eqs. [9]-[11]). In the most general case, $q = q(\Gamma_0)$, and the observational properties of the fireball would depend on the exact form of this dependence. We assume that q is only weakly dependent on Γ_0 , that is, we assume that q depends on Γ_0 through a function which varies much less strongly than $\propto \Gamma_0^{-4}$. In what follows, we examine the observational signatures of fireballs with different baryon loading Γ_0 but with constant q , keeping in mind the possibility of an underlying dependence of q on Γ_0 .

4. Prospects for Detecting Clean and Dirty Fireballs

The total number of photons collected from a GRB with energies near the mean photon energy of peak power output is roughly given by

$$N_\gamma \approx \frac{\Pi_0 t_d A_{\text{det}}}{4\pi d_L^2 m_e \mathcal{E}_0}, \quad (23)$$

where A_{det} is the effective area of the detector at energy $\mathcal{E}_0$. The deceleration time scale (5) defines the rough duration of peak power output, and eq. (10) gives the mean photon energy of the peak power output

at $t = t_d$. Hence

$$t_d(\text{s}) = 1730 \left(\frac{q}{\mathcal{E}_0} \right)^{2/3} [(1 - \eta/3)(1 + z)E_{54}]^{1/3} \cong 31 \left(\frac{q_{-3}}{\mathcal{E}_0} \right)^{2/3} [(1 - \eta/3)\zeta_6 E_{54}]^{1/3}, \quad (24)$$

where $q_{-3} = q/10^{-3}$ and $\zeta_6 \equiv (z + 1)/6$. Recalling the expression for the magnitude of peak-power output given by eq. (17), we obtain the result

$$N_\gamma \approx \frac{(2g - 3 + \eta)(3 - \eta)E_0}{2g(v^{-1} + \delta^{-1})(1 + z)} \frac{A_{\text{det}}}{4\pi m_e d_L^2 \mathcal{E}_0} \equiv K_\gamma \frac{E_{54} A_{\text{det}}}{d_{29}^2 \mathcal{E}_0 (1 + z)}, \quad (25)$$

where $d_{29} = d_L/(10^{29} \text{cm})$. Note that $d_{29} \cong 1$ when $z = 5$ in a $q_0 = 1/2$ cosmology with a Hubble constant equal to $65 \text{ km s}^{-1} \text{ Mpc}^{-1}$. For our standard parameters $\eta = 0$, $v = 4/3$, and $\delta = 0.2$,

$$K_\gamma = \begin{cases} 0.32, & \text{for } g = 1.6; \\ 2.4, & \text{for } g = 2.9. \end{cases} \quad (26)$$

4a. Clean ($\Gamma_0 \gg 300$) Fireballs

The 100 MeV threshold of EGRET corresponds to $\mathcal{E}_0 \simeq 200$. To produce 100 MeV nonthermal synchrotron photons mostly from electrons with energies $\simeq m_e \gamma_{e,\text{min}}$ requires fireballs with baryon-loading

$$\Gamma_0 \cong 76 \left[\frac{\mathcal{E}_0(1 + z)}{n_0^{1/2} q} \right]^{1/4} = 1420 \left(\frac{\zeta_6}{n_2^{1/2} q_{-3}} \right)^{1/4}, \quad (27)$$

where we set $\mathcal{E}_0 = 200$ on the right-hand-side of eq. (27). The number of photons detected from a clean fireball, given the standard parameters used in eqs. (25) and (26), is $N_\gamma \approx E_{54} A_{\text{det}} [\text{cm}^2] / [3750 d_{29}^2 \zeta_6]$ and $N_\gamma \approx E_{54} A_{\text{det}} [\text{cm}^2] / [490 d_{29}^2 \zeta_6]$ for $g = 1.6$ and $g = 2.9$, respectively. To detect clean fireballs requires that $N_\gamma > 1$. Thus sources of clean fireballs must be located at distances d_{29} given through the relation

$$\zeta_6^{1/2} d_{29} \lesssim E_{54}^{1/2} A_{1000}^{1/2} \begin{cases} 0.5 & \text{for } g = 1.6 \\ 1.4, & \text{for } g = 2.9 \end{cases}, \quad (28)$$

where $1000 A_{1000} \text{ cm}^2$ is the effective collecting area of the detector at $\mathcal{E}_0 \approx 200$.

The effective area of EGRET at 100 MeV in the wide field mode (e.g., Kurfess et al. 1997) is $\sim 800 \text{ cm}^2$ on-axis and $\sim 400 \text{ cm}^2$ at 20° off-axis, so clean fireballs would have to be located at $z \ll 5$ and at $z \lesssim 5$ to be detected with EGRET in the non-radiative and radiative limits, respectively. Bursts from clean fireballs with $\mathcal{E}_0 \simeq 200$ would have durations $t_d \approx 0.9 [q_{-3}^2 \zeta_6 (1 - \eta/3) E_{54}]^{1/3}$ (eq. [24]) at $z \cong 5$ and shorter durations when $z \ll 5$ and $q_{-3} \ll 1$. The spark chamber of EGRET requires $\approx 0.1 \text{ s}$ to recover from a photon event, although it can detect multiple photons if they arrive within a few microseconds. Consequently, EGRET is insensitive to more than one photon arriving between several microseconds and $\sim 100 \text{ ms}$, which is a critical range for the clean fireballs considered here.

It might be surprising that no sub-second bursts of gamma rays have been reported (Fichtel et al. 1994; Fichtel & Sreekumar 1997) from some members of the clean fireball class which are located at $z \lesssim 5$. We propose three possibilities to explain why this class of clean fireballs with peak power output near 100 MeV has not been discovered:

- There are many fewer clean fireballs, as defined by eq. (27), than those which produce the loaded BATSE GRBs.
- The duration of clean fireballs is typically $\lesssim 0.1$ s, so that deadtime limitations have prevented their discovery.
- Members of the clean fireball class evolve through cosmic time differently than the loaded fireballs which produce the GRBs detected with BATSE, and they do so in such a way to have escaped detection.

If the third point does not apply, then we can derive a requirement on the redshift z_B of the dimmest BATSE bursts to agree with the failure to detect clean fireballs with EGRET. Very crudely, we scale the number of clean fireballs which could be detected with EGRET to the BATSE detection rate ($\sim 800 \text{ yr}^{-1}$ full sky) and the EGRET field-of-view ($\sim 1/25$ th of the full sky) and lifetime (≈ 5 yrs). If clean fireballs can only be detected by EGRET within $z \lesssim z_E$ as given by eq. (28), then for EGRET not to have detected at least one member of this class, we find that

$$k_{cf} \times 0.04 \times \left(\frac{z_E}{z_B} \right)^j \times 800 \text{ yr}^{-1} \times 5 \text{ yr} \lesssim 1. \quad (29)$$

The term k_{cf} is a class enhancement factor correcting for the number of clean fireballs relative to the number of loaded fireballs which produce GRBs detectable with BATSE. The exponent j represents a cosmological scaling, and $j = 3$ for Euclidean space. Evolutionary and cosmic expansion effects will introduce modifications to j not considered here. Simply taking $j = 3$, we find that $z_E \lesssim 0.2 z_B k_{cf}^{-1/3}$ for the clean fireballs not to have been detected with EGRET. Noting eq. (28), this implies that even if $k_{cf} \sim 1$, sensitivity and FoV limitations could have prevented their discovery. Adding to the difficulty of the discovery is the deadtime limitations given by the second point above.

Stronger conclusions require a detailed size-distribution study involving the cosmological evolution of loaded GRB fireballs and clean fireballs in different scenarios, for example, the hypernova scenario (Paczynski 1998; Wijers et al. 1998). Given the uncertainty in the parameters, it remains an open question whether the larger effective area of the proposed *GLAST* instrument will be adequate for the discovery of the clean fireball class, or if all that is required is a wide FoV, large area gamma-ray calorimeter with fast timing. The discovery of subsecond bursts of ~ 100 MeV photons from clean fireballs remains a real possibility which, should such an event be detected, is not to be confused with the Hawking radiation from evaporating mini-black holes by virtue of its afterglow behavior.

Clean fireballs would be difficult to detect with the *Solar Maximum Mission*, which accumulated on time scales of 16 s and was not sensitive to short bursts with durations $\lesssim 2$ s (Harris & Share 1998). The short hard clean fireball bursts would also be difficult to detect with COMPTEL on *CGRO*, noting its smaller effective area ($\sim 10^2 \text{ cm}^2$; see, e.g., Hanlon et al. 1995). Nonetheless, further study of the COMPTEL sample of GRBs detected with BATSE, and a search for unidentified MeV transients will provide additional constraints on the number and redshift distribution of clean ($\Gamma_0 \gg 300$) fireballs.

4b. Dirty ($\Gamma_0 \ll 300$) Fireballs

Because a large number of X-ray telescopes operate in the 1-10 keV range, we can use $\mathcal{E}_0 \sim 3/511$ to define the dirty fireball regime, which includes those blast waves with

$$\Gamma_0 \simeq 21 \left[\frac{(1+z)}{n_0^{1/2} q} \right]^{1/4} \simeq 104 \left(\frac{\zeta_6}{n_2^{1/2} q_{-3}} \right)^{1/4} \quad (30)$$

(compare eq. [27]). Detection of dirty fireballs with X-ray telescopes is difficult because of the intense backgrounds. A signal of N_γ counts given by eq. (23) is detected above a background of B_G counts at the n_σ significance level provided that

$$N_\gamma \gtrsim n_\sigma (2B_G)^{1/2}. \quad (31)$$

The number of background counts

$$B_G \cong \Delta \epsilon_{\text{det}} \cdot \Phi_{\text{dif}}(\Omega, \epsilon_{\text{det}}) \cdot \Delta \Omega \cdot \Delta t \cdot A_{\text{det}}, \quad (32)$$

where, from right to left, the terms are the detector area, the observing time, the field-of-view (FoV) of the detector, the diffuse photon flux weighted by an average photon energy within the band, and the bandwidth. The background noise is composed of instrumental, magnetospheric, zodiacal, Galactic, and extragalactic components, which are time-dependent in general. Here we give a background estimate for X-ray telescopes considering only the diffuse cosmic X-ray noise background, yielding a lower limit to B_G .

In the 3-100 keV range, Boldt (1987) gives the expression

$$\epsilon \Phi_{\text{XRB}}(\epsilon, \Omega) \left[\frac{\text{ph}}{\text{cm}^2 \text{ s sr}} \right] \simeq 5.6 \left[\frac{E(\text{keV})}{3 (\text{keV})} \right]^{-0.29} \exp[-E(\text{keV})/40 \text{ keV}] \quad (33)$$

for the diffuse extragalactic X-ray background flux. At 3 keV, corresponding roughly to the mean energy of photons detected by an X-ray telescope sensitive in the 1-10 keV band, the diffuse flux is $\simeq 5 \text{ ph cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$.

Substituting eqs. (23) and (32) into eq. (31) with the stated diffuse flux gives the result that detection at the n_σ level requires an X-ray telescope with

$$\frac{A_{\text{det}}}{\Delta \Omega} \left[\frac{\text{cm}^2}{\text{sr}} \right] \gtrsim \frac{3.4 \times 10^{-4} d_{29}^4 n_\sigma^2 (1+z)^2 \Delta t}{K_\gamma^2 E_{54}^2 f(\Delta t/t_d)} \quad (34)$$

The function $f(y)$ weights the significance of detection by the observing interval over the period in which the fireball's emissions are monitored. Its form depends on the time-dependence of the light curve of the fireball emissions, which can be modeled using the eq. (1), and the epoch of observation. In an illustrative regime, $f(y) = y^2$ for $y \leq 1$, and $f(y) = 1$ for $y > 1$. The detection efficiency is clearly optimized when $\Delta t \approx t_d$. The deceleration time scale for a burst peaking at $\mathcal{E}_0 \cong 3/511 = 0.006$ is

$$t_d(\text{s}) \simeq 970 [\zeta_6 q_{-3}^2 (1 - \eta/3) E_{54}]^{1/3}. \quad (35)$$

The search for dirty fireballs should be conducted in the range of time scales from 10^2 - 10^4 s, recognizing the difficulty due to orbital variations of the background radiation environment.

When $\eta = 0$, one obtains the requirement for an X-ray telescope to be able to detect a dirty fireball at the n_σ level, namely

$$\frac{A_{\text{det}}}{\Delta \Omega} \gtrsim \frac{\zeta_6^{7/3} d_{29}^4 (n_\sigma/5)^2 q_{-3}^{2/3}}{E_{54}^{5/3}} \begin{cases} 940, & \text{for } g = 1.6; \\ 125, & \text{for } g = 2.9. \end{cases} \quad (36)$$

Typical X-ray detectors have effective areas $\sim 10^2 \text{ cm}^2$ (see Table 3 for general specifications of some X-ray telescopes used to detect GRBs), so we scale to a detector of area $100A_{100} \text{ cm}^2$. To detect dirty fireballs above the diffuse X-ray background means that the detector’s FoV must satisfy the relation

$$\frac{\Delta\Omega}{4\pi} \lesssim \frac{E_{54}^{5/3} A_{100}}{\zeta_6^{7/3} d_{29}^4 q_{-3}^{2/3} (n_\sigma/5)^2} \begin{cases} 0.0027, & \text{for } g = 1.6; \\ 0.15, & \text{for } g = 2.9. \end{cases} \quad (37)$$

Thus an X-ray detector has to have a FoV $\lesssim 10^\circ \times 10^\circ$ or $\lesssim 80^\circ \times 80^\circ$ in order to detect dirty fireballs at $z \sim 5$, depending on whether the blast wave decelerates primarily in the non-radiative or radiative regimes, respectively. Fireballs at $z \ll 1$ place much weaker constraints on the detector’s FoV, as indicated in equation (37), and then it becomes a question of the number of such sources one could expect to see.

Small FoV instruments will only detect an X-ray flash from a dirty fireball very rarely unless the class of dirty fireballs is populous. Given that there are ~ 800 BATSE-detectable GRBs per year averaged over the full sky, this implies a chance probability of $\sim 0.02 \cdot k_{df} \cdot k_d^2$ per year for a detector with a $k_d^\circ \times k_d^\circ$ FoV. The class enhancement factor k_{df} of dirty fireballs compared to loaded BATSE/GRB fireballs is not known at present, though it can be strongly constrained by modeling the duration distribution of GRBs detected with BATSE or with other burst detectors (Dermer et al. 1998). Searches for dirty fireballs should be made with the Wide Field Camera data on Beppo-SAX, and with future wide FoV imaging X-ray detectors. Concrete deductions about the probability of detecting serendipitous dirty fireball X-ray flashes requires a size distribution study tied to a particular detector’s characteristics.

It is feasible that some of the X-ray flashes observed in archival searches of HEAO (Connors, Serlemitsos, & Swank 1986) and *ROSAT* PSPC (Li et al. 1998; Sun et al. 1998) data are emissions from dirty fireballs only if $k_{df} \gg 1$, given that the effective FoV of the *ROSAT* PSPC is $\sim 1^\circ \times 1^\circ$. The GRBs detected with *Ginga* (Strohmayer et al. 1998) have νF_ν peaks which are on average at lower energies than those detected with BATSE. Because *Ginga* was sensitive to lower energies events than BATSE, this might indicate that it is more sensitive to a population of GRBs with greater baryon-loading. A straightforward prediction from this model is that the mean duration of detected GRBs is inversely correlated with the triggering energy range.

5. Discussion

A slewing strategy such as the one so successfully demonstrated by the Beppo-SAX team and collaborators (e.g. Costa et al. 1997; van Paradijs et al. 1997; Piro et al. 1998a,b) opens rich possibilities for discoveries which must be carefully weighed in GRB telescope design. Here we have attempted to provide an exposition of the blast-wave model which provides a simple way to model the evolving spectral power fluxes, and which furthermore helps us to understand dedicated and serendipitous GRB observations.

Eq. (1) is the proposed time-dependent spectral form. It employs the 9 parameters listed in Table 2 that follow from the basic blast-wave model. Its crucial underlying assumptions are that the photon indices ν and δ and the equipartition term q are constant in time, and that q is independent or only weakly dependent on Γ_0 . The energies of photons which carry the bulk of the power are $\propto q\Gamma_0^4$ and are therefore most strongly dependent on the value of Γ_0 . Over this looms a question of lack of self-consistency, since specification of the radiative regime g constrains the magnetic field and photon energy densities and particle distributions to produce the requisite radiative power. Comparison of eq. (1) with the results of a numerical code employing a self-consistent treatment of plasmoid dynamics (Chiang & Dermer 1998) indicates that

the proposed analytic description of the blast wave emission spectra remains accurate except when the blast wave is highly radiative.

In spite of these concerns, eq. (1) can be easily employed to make estimations of imaging capabilities of burst detectors given slewing rates of narrow field instruments toward a cosmic transient. It can be used to calculate photoelectric absorption variations in the GRB environment and its imprint upon the observed burst spectrum (Böttcher et al. 1998). It can also, of course, be used to model the temporal behavior of GRB spectra and to examine GRB phenomenology in terms of the fireball/blast-wave model (Dermer et al. 1998).

By displaying the spectral power flux in the form of eq. (1), we obtained an important rule on fireball emissions relating the observed peak power Π_0 , the characteristic duration t_d of this peak power output, and the mean energy $\mathcal{E}_0$ of photons which carry this energy. The characteristic time scale over which the bulk of the power is radiated is the deceleration time scale t_d (eq. [5]; Rees & Mészáros 1992), and $t_d \propto \Gamma_0^{-8/3}$, where Γ_0 specifies the baryon loading. The peak power output Π_0 (eq. [17]) is also $\propto \Gamma_0^{8/3}$, assuming that the bulk of the dissipated energy is in the form of photon radiation. Because most of the total particle kinetic energy, E_0 , originally in the fireball during its coasting phase is radiated during the observed time t_d , we have

$$E_0 \approx \Pi_0 t_d . \quad (38)$$

Precisely,

$$\Pi_0 t_d = \frac{(2g - 3 + \eta)(3 - \eta)}{2g(v^{-1} + \delta^{-1})(1 + z)} E_0 = \begin{cases} 0.033 E_0 / (1 + z), & \text{for } g = 1.6; \\ 0.25 E_0 / (1 + z), & \text{for } g = 2.9. \end{cases} \quad (39)$$

(In the evaluation on the right-hand-side of this expression, we let $\eta = 0$, $v = 4/3$, and $\delta = 0.2$.) Moreover, the bulk of this energy is observed in the form of photons with characteristic energy $\mathcal{E}_0$ given by eq. (10); note that $\mathcal{E}_0 \propto q\Gamma_0^4$. For these parameters, a full 25% of the energy of the GRB source is radiated within the deceleration time scale for a highly radiative fireball.

Eqs. (38) and (39) can be used to summarize the emission properties of fireballs with different baryon loading and prospects for detecting such fireballs. Fig. 3 shows the variation of Π_0 , t_d , and $\mathcal{E}_0$ when Γ_0 ranges from 100 to 1000. The dependence of these observable due to variations in E_{54} and n_0 are also indicated. The overall powers, time scales, and photon energies of peak power output of fireballs with $\Gamma_0 \sim 300$, and for clean and dirty fireballs can be determined from this figure.

We used benchmark characteristics of X-ray, soft gamma-ray, and medium-energy gamma-ray telescopes to establish criteria for clean ($\Gamma_0 \gg 300$), loaded ($\Gamma_0 \sim 300$), and dirty ($\Gamma_0 \ll 300$) fireballs. Detection of clean fireballs at $\gtrsim 100$ MeV energies implies $\Gamma_0 \gtrsim 1000$ (eq. [27]), and the detection of dirty fireballs at ~ 3 keV energies implies $\Gamma_0 \lesssim 100$ (eq. [30]). The clean fireball class was not discovered with EGRET; this means either that there are relatively few sources of energetic clean fireballs, or that the powers and durations of clean fireballs are such that sensitivity and deadtime limitations of previous > 100 MeV γ -ray detectors have prevented their detection.

Here we qualify our use of eq. (38) which gives the general observational properties of fireballs with different baryon loading parameters, EGRET has, of course, detected > 100 MeV gamma-ray emission from GRBs detectable with BATSE from the loaded ($100 \lesssim \Gamma_0 \lesssim 1000$) fireball class. Many of these sorts of GRBs will undoubtedly be detected with the *GLAST* mission as well. But the > 100 MeV and GeV photons detected to date (e.g., Hurley et al. 1994; Dingus 1995) are not those produced by the electrons near the low-energy cutoff of the electron distribution function. These photons are much more likely to be

the SSC emission (e.g., Chiang & Dermer 1998) or the nonthermal synchrotron radiation from ultra-high energy protons (Vietri 1997b; Böttcher & Dermer 1998) accelerated in the loaded fireball blast wave. These emissions are distinguished by their slow decay, which explains why EGRET often detects this emission long after the main portion of the BATSE burst has decayed. What EGRET has not discovered but is implied by the fireball/blast-wave model is a class of clean fireballs which produces a luminous subsecond burst of $\gg 100$ MeV radiation. The prediction of such a discovery with improved γ -ray telescopes is made here.

We also used analyses outlined by eq. (38) to describe the properties of dirty fireballs. Because Π_0 and $\mathcal{E}_0$ are lower in dirtier fireballs, the diffuse sky backgrounds make detection of such a class of objects by wide-field instruments problematic. On the other hand, the expected rate of detection of dirty fireballs by pointed instruments with a few square degree FoV is very unlikely unless the rate of dirty fireball explosions greatly exceeds the rate of loaded BATSE/GRB fireballs. Whether any members of this class has been discovered is uncertain, though X-ray flashes have indeed been detected in the *HEAO* and *ROSAT* data bases. We predict that moderate FoV instruments ($\sim 1\%$ of the full sky) with surveying capabilities will discover dirty fireball X-ray transients with typical durations of 10^2 - 10^4 seconds.

Beppo-SAX has shown that improved imaging and rapid response are key to making progress in solving the GRB mystery. Granted this success, the design of a new GRB mission should therefore employ a hard X-ray/soft GRB detector and a narrow FoV X-ray telescope which can rapidly slew to the GRB position. But to discover the sister classes of clean and dirty fireballs implied by the fireball/blast-wave model will require additional X-ray and gamma-ray capabilities as outlined above and considered in more detail elsewhere (Dermer et al. 1998).

The work of CD was supported by the Office of Naval Research and the *Compton Gamma Ray Observatory* Guest Investigator Program. CD thanks Jim Kurfess, Neil Johnson, Mark Strickman, and Gerry Share for enlightening discussions, and the referee for a useful critique. The work of JC was performed while he held a National Research Council - Naval Research Laboratory Associateship. MB acknowledges support by the German Academic Exchange Service (DAAD).

Appendix A. The Fraction of Radiated Energy

An expression is derived in the asymptotic regime $x \gg x_d$ which relates the fraction ζ of swept-up energy retained in the blast wave to the index g which characterizes the radiative regime and evolution of the Lorentz factor Γ time scale of the blast wave (see eq. [4]). We start from the equation of momentum conservation (eq. [6] of Dermer & Chiang [1998]) and note that the energy in swept-up particles accumulated up to the point x is equal to

$$\int_0^\infty dp \gamma N_p(\gamma) = \int_0^x d\tilde{x} n_{\text{ext}}(\tilde{x}) A(\tilde{x}) [\zeta \Gamma(\tilde{x}) + (1 - \zeta)]. \quad (\text{A1})$$

In the asymptotic regime $x_d \ll x \ll x_{\text{tr}}$, where $x_{\text{tr}} = x_d \Gamma_0^{1/g}$ and $\Gamma(x) = \Gamma_0 (x/x_d)^{-g} \gg 1$, this yields

$$-\frac{\Gamma'(x)}{\Gamma^2(x)} = \frac{g}{x \Gamma(x)} \approx \frac{n_0 A_0 \left(\frac{x}{x_d}\right)^{2-\eta}}{N_{\text{th}} + n_0 A_0 \left[\Gamma_0 \zeta \int_{x_d}^x d\tilde{x} \left(\frac{\tilde{x}}{x_d}\right)^{2-g-\eta} + (1 - \zeta) \int_0^x d\tilde{x} \left(\frac{\tilde{x}}{x_d}\right)^{2-\eta} \right]}. \quad (\text{A2})$$

Unless the blast wave is in the extreme radiative regime, the denominator on the right-hand side of eq. (A2) is dominated by the first integral. Thus if $x \ll \zeta x_{\text{tr}}$, eq. (A2) becomes

$$\frac{g}{\Gamma_0 x_d} \left(\frac{x}{x_d} \right)^{g-1} \approx \frac{3-g-\eta}{\zeta \Gamma_0 x_d} \left(\frac{x}{x_d} \right)^{g-1}, \quad (\text{A3})$$

or

$$\zeta = \frac{3-g-\eta}{g}. \quad (\text{A4})$$

Table 1: Slopes of the Temporal Evolution of the Flux at Fixed Photon Energy when $t \gg t_d$ and $\eta = 0$

slope	g	$v = 1$	$v = 4/3$					
χ_1	1.6	0.48	0.98					
χ_1	2.9	0.29	0.86					
		$\delta = 0$	$\delta = 1/5$	$\delta = 1/3$	$\delta = 1/2$	$\delta = 3/4$	$\delta = 1$	$\delta = 2$
χ_2	1.6	1.05	1.35	1.56	1.81	2.19	2.57	4.09
χ_2	2.9	1.41	1.75	1.98	2.26	2.69	3.12	4.82

Table 2: Standard Parameters for Model of Evolving GRB Spectral Power Flux

Parameter	Standard Value	Description
Γ_0	300	initial bulk Lorentz factor
q	10^{-3}	equipartition term
g	2	index of Γ evolution
E_0	10^{54} ergs	total particle kinetic energy in GRB
v	$4/3$	spectral index of rising portion of νL_ν spectrum
δ	0.2	spectral index of falling portion of νL_ν spectrum
z	5	cosmological redshift
n_0	10^2 cm^{-3}	density at $x = x_d$
η	0	index of density distribution

Table 3: Characteristics of Operating Missions with GRB Detectors

	Energy band (keV)	FoV	Effective Area (cm ²)	Sensitivity ^b	Imaging
CGRO BATSE	50 – 300	4 π	1800 cm ² ^a	6 $\times$ 10 ⁻⁸ ^c	few degrees
SAX GRBM	40 – 700	20° $\times$ 20°	120		10' $\rightarrow$ 3'
SAX WFC	1.5 – 26	20° $\times$ 20°	120	$\sim$ 10 ⁻¹⁰ in 10 ³ s ^d	10' $\rightarrow$ 2'
SAX NFI LECS	0.1 – 10	0.5°	22@0.25 keV	$\sim$ 3 $\times$ 10 ⁻¹⁴ in 10 ⁵ s	50''
SAX NFI MECS	0.1 – 10	0.5°	150@6 keV		

^aEach Large Area Detector

^bergs cm⁻² s⁻¹

^cLarge Area Detectors; 10 s GRB

^dBeppo-SAX data from homepage; see also Piro et al. (1998)

REFERENCES

- Band, D. L., et al. 1993, ApJ, 413, 281
- Belli, M. 1995, Ap&SS, 231, 43
- Blandford, R. D., & McKee, C. F. 1976, Phys. Fluids, 19,1130
- Boldt, E. 1987, Physics Reports, 146, no. 4, 215
- Böttcher, M., & Dermer, C. D. 1998, ApJ, 499, L131
- Böttcher, M., Crider, A. W., Liang, E. P., & Dermer, C. D. 1998, in preparation
- Catelli, J. R., Dingus, B. L., & Schneid, E. J. 1996, in the Third Huntsville Symposium on Gamma Ray Bursts, ed. C. Kouveliotou, M. F. Briggs, & G. J. Fishman (New York: AIP), 158
- Cavallo, G., & Rees, M. J. 1978, 183, 359
- Chiang, J., & Dermer, C. D. 1998, ApJ, submitted (astro-ph/9803339)
- Chiang, J. 1998, ApJ, in press (astro-ph/9805302)
- Cohen, E., Katz, J. I., Piran, T., Sari, R., Preece, R. D., & Band, D. L. 1997, ApJ, 488, 330
- Connors, A., Serlemitsos, P., & Swank, J. H. 1986, ApJ, 303, 769
- Costa, E., et al. 1997, Nature, 387, 783
- Dermer, C. D., Böttcher, M., & Chiang, J. 1998, in preparation
- Dermer, C. D., & Chiang, J. 1998, New Astronomy, 3, 157
- Dermer, C. D., & Mitman, K. E. 1998, ApJL, submitted
- Dermer, C. D. 1998, ApJ, 501, L157
- Dermer, C. D., Sturmer, S. J., & Schlickeiser, R. 1997, ApJS, 109, 103
- Dingus, B. L. 1995, A&SS, 231, 187

- Djorgovski, S. G. 1997, *Nature*, 387, 876
- Fenimore, E. E., In't Zand, J. J. M., Norris, J. P., Bonnell, J. T., & Nemiroff, R. J. 1995, *ApJ*, 448, L101
- Feroci, M. 1998, *A&A*, 323, L29
- Fichtel, C. E., & Sreekumar, P. 1997, in *The Fourth Compton Symposium*, ed. C. D. Dermer, M. S. Strickman, & J. D. Kurfess (New York: AIP), 436
- Fichtel, C. E., et al. 1994, *ApJ*, 434, 557
- Frail, D. 1998, in the *Fourth Huntsville Gamma-Ray Burst Symposium*, ed. C. A. Meegan, R. D. Preece, & T. M. Koshut (AIP: New York), 563
- Galama, T. J., et al. 1998, *ApJ*, 497, L13
- Groot, P. J., et al. 1998, *ApJ*, 502, L123
- Harris, M. J., & Share, G. J. 1998, *ApJ*, 494, 724
- Hanlon, L. et al., 1995, *A&SS*, 231, 157
- Hurley, K. C., et al. 1994, *Nature*, 372, 652
- Katz, J. I. 1994, *ApJ*, 432, L107
- Kouveliotou, C., Meegan, C. A., Fishman, G. J., Bhat, N. P., Briggs, M. S., Koshut. T. M., Paciesas, W. S., & Pendleton, G. N. 1993, *ApJ*, 413, L101
- Kulkarni, S. et al. 1998a, *Nature*, 393, 35
- Kulkarni, S. et al. 1998b, *Nature*, submitted (astro-ph/9807001)
- Kurfess, J. D., Bertsch, D. L., Fishman, G. J., & Schönfelder, V. 1997, in *The Fourth Compton Symposium*, ed. C. D. Dermer, M. S. Strickman, & J. D. Kurfess (New York: AIP), 509
- Li, H., Sun, X., Fenimore, E. E., & Wang, Q. D., in the *Fourth Huntsville Gamma-Ray Burst Symposium*, C. A. Meegan, R. D. Preece, & T. M. Koshut, eds. (New York: AIP), 420
- Lozinskaya, T. A. 1992, *Supernovae and Stellar Winds in the Interstellar Medium* (New York: AIP)
- Marscher, A. P., & Gear, W. K. 1985, *ApJ*, 298, 114
- Mészáros, P., & Rees, M. J. 1993a, *ApJ*, 418, L59
- Mészáros, P., & Rees, M. J. 1993b, *ApJ*, 405, 278
- Mészáros, P., & Rees, M. J. 1997, *ApJ*, 476, 232
- Mészáros, P., Rees, M. J., & Papathanassiou, H. 1994, *ApJ*, 432, 181
- Mészáros, P., Rees, M. J., & Wijers, R. A. M. J., 1997, *ApJ*, submitted (astro-ph/9709273)
- Mészáros, P., Laguna, P., & Rees, M. J. 1993, *ApJ*, 415, 181
- Metzger, M. R., et al. 1997, *Nature*, 387, 878
- Paczynski, B., & Rhoads, J. 1993, *ApJ*, 418, L5
- Paczynski, B. 1998, *ApJ*, L45
- Panaitescu, A., & Mészáros, P. 1998a, *ApJ*, 493, L31
- Panaitescu, A., & Mészáros, P. 1998b, *ApJ*, 492, 683
- Panaitescu, A., & Mészáros, P. 1998c, *ApJ*, 501, 772
- Piro, L., et al. 1998a, *A&A*, 331, L41

- Piro, L., et al. 1998b, *A&A*, 329, 906
- Piran, T., & Shemi, A. 1993, *ApJ*, 403, L67
- Rees, M. J., & Mészáros, P., 1992, *MNRAS*, 258, 41P
- Ryu, D., & Vishniac, E. 1991, *ApJ*, 368, 411
- Sahu, K. C. et al. 1997, *Nature*, 387, 476
- Sari, R. 1997, *ApJ*, 489, L37
- Sari, R. 1998, *ApJ*, 494, L49
- Sari, R., Piran, T., & Narayan, R. 1998, 497, L17
- Shemi, A., & Piran, T. 1990, *ApJ*, 365, L55
- Strohmayer, T. E., Fenimore, E. E., Murakami, T., & Yoshida, A. *ApJ*, 1998, 500, 873
- Sun, X., Li, H., Fenimore, E. E., & Wang, Q. D., in the Fourth Huntsville Gamma-Ray Burst Symposium, C. A. Meegan, R. D. Preece, & T. M. Koshut, eds. (New York: AIP), 456
- Tavani, M. 1996, *Phys. Rev. Letters*, 76, 3478
- van Paradijs, J., et al. 1997, *Nature*, 386, 686
- Vietri, M. 1997a, *ApJ*, 478, L9
- Vietri, M. 1997b, *PRL* 78, 23, 4328
- Waxman, E. 1997a, *ApJ*, 485, L5
- Waxman, E. 1997b, *ApJ*, 491, L19
- Wijers, R. A. M. J., Bloom, J. S., Bagla, J. S., & Natarajan, P. 1998, *MNRAS*, 294, L13
- Wijers, R. A. M. J., Mészáros, P., & Rees, M. J. 1997, *MNRAS*, 288, L51




Fig. 1.— Spectra and light curves for a model GRB using the standard parameters in Table 2. (a) Evolution of model GRB spectra at different observing times from $1 \mu\text{s}$ to 10^7 s are shown in factor-of-10 increments. Solid lines plot the trajectory of the peak power $P_p(t)$ as a function of the photon energy $\epsilon_p(t)$ of the peak of the νL_ν spectrum at different observing times. (b) GRB light curves at different observing energies for the model GRB spectra shown in (a).




Fig. 2.— Trajectories of model GRBs in the $P_p(t)$ - $\epsilon_p(t)$ plane. All parameters are given in Table 2, except as noted. (a) Trajectories for different values of the index g characterizing the radiative regime and evolution of Γ through eq. (4).




Fig. 2.— (b) Same as Fig. 2a, but for different values of the density n_0 of the circumburst medium.




Fig. 2.— (c) Same as Fig. 2a, but for different values of the total explosion energy E_0 .




Fig. 2.— (d) Same as Fig. 2a, but for different values of the equipartition factor q .




Fig. 2.— (e) Same as Fig. 2a, but for different values of the baryon-loading factor Γ_0 . Note the strong dependence of the duration, peak power, and photon energy of the νL_ν peak on Γ_0 .




Fig. 3.— Dependence of the observables Π_0 , t_d and $\mathcal{E}_0$ on the baryon-loading parameter Γ_0 , the total explosion energy $E_0 = 10^{54}E_{54}$ ergs and the external density n_0 (cm^{-3}), for Γ_0 ranging from 100 to 1000. Other parameters are given in Table 2.